


RYBREVANT® (amivantamab-vmjw) is given through an intravenous infusion under the attention of your healthcare team

- Your healthcare provider will decide the time between doses as well as how many treatments you will receive
- Your healthcare provider will decide the appropriate dose of RYBREVANT® based on your body weight

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins and herbal supplements

Infusion Schedule

With RYBREVANT®, the dosing schedule may change and become less frequent after your first dose.



While everyone's experience is unique, infusion times may get shorter after your first dose.

- Without an infusion-related reaction (IRR):
 - Your first dose (the infusion on Day 1 and Day 2) can take about 4 to 6 hours
 - Infusions will get shorter starting on Week 2, and can take about 2 to 4 hours

Your healthcare provider will decide:

- The time between doses
- How many treatments you will receive
- The appropriate dose of RYBREVANT® based on your body weight

However, your healthcare provider may interrupt or stop your infusion if an adverse reaction occurs.

What is RYBREVANT® (amivantamab-vmjw)?

RYBREVANT® is a prescription medicine used to treat adults with non-small cell lung cancer (NSCLC) that:

- has spread to other parts of the body (metastatic) or cannot be removed by surgery, **and**
- has a certain abnormal epidermal growth factor receptor "EGFR" gene(s) **and**
- whose disease has worsened while on or after chemotherapy that contains platinum.

Your healthcare provider will perform a test to make sure that RYBREVANT® is right for you.

It is not known if RYBREVANT® is safe and effective in children.

RYBREVANT® is approved based on medical studies that measured how many patients responded to treatment. There are ongoing studies to confirm the continued approval of RYBREVANT®.

IMPORTANT SAFETY INFORMATION

What are the possible side effects of RYBREVANT®?

RYBREVANT® may cause serious side effects including infusion-related reactions, lung problems, skin problems, and eye problems.

Please read Important Safety Information throughout and on pages 4-5.

Please read enclosed full Prescribing Information for RYBREVANT®.

Preparing for infusions

There are a few tips to keep in mind to ensure you're ready for an infusion:



It may help to drink plenty of water the day before an infusion and have a healthy breakfast the morning of. Consider bringing a snack and drink with you to your appointment, but check with the facility first.



Bring a puzzle, game, or book—something to help pass the time during your infusion.



Infusions can last several hours, so make sure you've set aside enough time for your appointment.



Pay close attention to how you feel during an infusion, and make sure you let your healthcare team know if you experience any discomfort—they know how to help.



Consider arranging transportation to and from the infusion center. You may receive medicines before your infusion that can make you drowsy. If you want to make these arrangements and need help doing so, reach out to your support network.



Feeling unsure about your infusion? Call your healthcare team. They can help to answer any questions that you may have along the way.



Loose clothing can help keep you comfortable in the infusion room and allow your healthcare team to access the infusion site easily. Bring a pillow and blanket for extra comfort.

Important Safety Information

What is RYBREVANT® (amivantamab-vmjw)?

RYBREVANT® is a prescription medicine used to treat adults with non-small cell lung cancer (NSCLC) that:

- has spread to other parts of the body (metastatic) or cannot be removed by surgery, **and**
- has a certain abnormal epidermal growth factor receptor “EGFR” gene(s) **and**
- whose disease has worsened while on or after chemotherapy that contains platinum.

Your healthcare provider will perform a test to make sure that RYBREVANT® is right for you.

It is not known if RYBREVANT® is safe and effective in children.

RYBREVANT® is approved based on medical studies that measured how many patients responded to treatment. There are ongoing studies to confirm the continued approval of RYBREVANT®.

IMPORTANT SAFETY INFORMATION

Before you receive RYBREVANT®, tell your healthcare provider about all of your medical conditions, including if you:

- have a history of lung or breathing problems
- are pregnant or plan to become pregnant. RYBREVANT® can harm your unborn baby.

Females who are able to become pregnant:

- Your healthcare provider should do a pregnancy test before you start treatment with RYBREVANT®.
 - You should use effective birth control (contraception) during treatment and for 3 months after your final dose of RYBREVANT®.
 - Tell your healthcare provider right away if you become pregnant or think you might be pregnant during treatment with RYBREVANT®.
- are breastfeeding or plan to breastfeed. It is not known if RYBREVANT® passes into your breast milk. Do not breastfeed during treatment and for 3 months after your final dose of RYBREVANT®.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

How will I receive RYBREVANT®?

- RYBREVANT® will be given to you by your healthcare provider by intravenous infusion into your vein.
- Your healthcare provider will decide the time between doses as well as how many treatments you will receive.
- Your healthcare provider will give you medicines before each dose of RYBREVANT® to help reduce the risk of infusion-related reactions.
- If you miss any appointments, call your healthcare provider as soon as possible to reschedule your appointment.

What should I avoid while receiving RYBREVANT®?

RYBREVANT® can cause skin reactions. You should limit your time in the sun during and for 2 months after your treatment with RYBREVANT®. Wear protective clothing and use sunscreen during treatment with RYBREVANT®.

What are the possible side effects of RYBREVANT®?

RYBREVANT® may cause serious side effects, including:

- **infusion-related reactions.** Infusion-related reactions are common with RYBREVANT® and can be severe or serious. Tell your healthcare provider right away if you get any of the following symptoms during your infusion of RYBREVANT®:
 - shortness of breath
 - fever
 - chills
 - nausea
 - flushing
 - chest discomfort
 - lightheadedness
 - vomiting

- **lung problems.** RYBREVANT® may cause lung problems that may lead to death. Symptoms may be similar to those symptoms from lung cancer. Tell your healthcare provider right away if you get any new or worsening lung symptoms, including shortness of breath, cough, or fever.
- **skin problems.** RYBREVANT® may cause rash, itching, and dry skin. You may use alcohol-free moisturizing cream for dry skin. Tell your healthcare provider right away if you get any skin reactions. Your healthcare provider may treat you with a medicine(s) or send you to see a skin specialist (dermatologist) if you get skin reactions during treatment with RYBREVANT®. See “What should I avoid while receiving RYBREVANT®?”
- **eye problems.** RYBREVANT® may cause eye problems. Tell your healthcare provider right away if you get symptoms of eye problems which may include:
 - eye pain
 - dry eyes
 - eye redness
 - blurred vision
 - changes in vision
 - itchy eyes
 - excessive tearing
 - sensitivity to light

Your healthcare provider may send you to see an eye specialist (ophthalmologist) if you get eye problems during treatment with RYBREVANT®. You should not use contact lenses until your eye symptoms are checked by a healthcare provider.

The most common side effects of RYBREVANT® include:

- rash
- infusion-related reactions
- infected skin around the nail
- muscle and joint pain
- shortness of breath
- nausea
- feeling very tired
- swelling of hands, ankles, feet, face, or all of your body
- sores in the mouth
- cough
- constipation
- vomiting
- changes in certain blood tests

Your healthcare provider may temporarily stop, decrease your dose or completely stop your treatment with RYBREVANT® if you have serious side effects.

These are not all of the possible side effects of RYBREVANT®.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

General information about safe and effective use of RYBREVANT®

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. You can ask your healthcare provider or pharmacist for information about RYBREVANT® that is written for health professionals.

Please read enclosed full Prescribing Information for RYBREVANT®.

cp-213277v2

What you should know about infusion-related reactions (IRRs)

Your healthcare team is trained to help you with any IRRs you may experience, so make sure to tell them how you're feeling during your infusion

Here are some additional important things you should know about IRRs:

- Infusion-related reactions are common with RYBREVANT® (amivantamab-vmjw) and can be severe or serious
- IRRs can involve shortness of breath, fever, chills, nausea, flushing, chest discomfort, lightheadedness, or vomiting that occur during your infusion
- IRRs were one of the most common side effects in a clinical study, and occurred in 66% of people treated with RYBREVANT®
- 65% of IRRs occur on Day 1 of the first dose, and typically happen within the first hour of an infusion. On Day 2 of the first dose, 3.4% of people experienced an IRR. During Week 2, 0.4% of people experienced an IRR. During all other infusions, 1.1% of people experienced an IRR
- Your healthcare provider will give you medicines before each dose of RYBREVANT® to help reduce the risk of IRRs

Pay close attention to how you feel during an infusion, and make sure you let your healthcare team know if you experience any discomfort—they know how to help



IMPORTANT SAFETY INFORMATION (CONTINUED)

What are the possible side effects of RYBREVANT®?

RYBREVANT® may cause serious side effects, including:

- **infusion-related reactions.** Infusion-related reactions are common with RYBREVANT® and can be severe or serious. Tell your healthcare provider right away if you get any of the following symptoms during your infusion of RYBREVANT®:
 - shortness of breath
 - flushing
 - fever
 - chest discomfort
 - chills
 - lightheadedness
 - nausea
 - vomiting

Please read Important Safety Information throughout and on pages 4-5.
Please read enclosed full Prescribing Information for RYBREVANT®.

